

# Pulmonary alveolar microlithiasis: review of Turkish reports

E S Ucan, A I Keyf, R Aydılek, Z Yalcin, S Sebit, M Kudu, U Ok

Gülhane Military Medical Faculty, Haydarpaşa Training Hospital

Department of Chest Diseases, Medical School, Dokuz Eylül University, Izmir, Turkey

E S Ucan  
A I Keyf  
R Aydılek  
Z Yalcin  
S Sebit  
M Kudu  
U Ok

Reprint requests to:  
Dr E S Ucan,  
Türkiye Cad.No 33,  
Bornova, Izmir, Turkey

Received 28 February 1992  
Returned to authors  
1 April 1992  
Revised version received  
26 May 1992

Accepted 9 July 1992

## Abstract

**Pulmonary alveolar microlithiasis is a rare disorder, only 173 cases having been reported worldwide. Fifty two cases from Turkey are reported, 49 of which have previously been described only in Turkish publications. The mean age of the patients was 27 (SD 12) years, 34 were male, and 10 were symptomless. In 40 of the 52 cases diagnosis was confirmed histopathologically. Nineteen cases were diagnosed in siblings. This high rate suggests that pulmonary alveolar microlithiasis is a familial disease, which, though rare, is for unknown reasons most common in Turkey.**

(Thorax 1993;48:171-173)

Pulmonary alveolar microlithiasis is a rare idiopathic disease characterised by microliths in the alveoli, first described by Friedrich in 1856<sup>1</sup> and then by Harbitz in 1918.<sup>2</sup> O'Neill reported a family with pulmonary alveolar

microlithiasis in 1967 and noted that the total number of recorded cases was 70.<sup>3,4</sup> Subsequently the number of published cases has risen to 173.<sup>5-7</sup> The first Turkish case was reported in 1962, and the total number of Turkish cases reached 52 this year.<sup>8</sup> As none of the Turkish cases, however, were included in any previous series we reviewed them and summarise their features here.<sup>3,4,6,7</sup>

## Cases

In all there were 34 male and 18 female patients with a mean age of 27.3 (SD 12) (range 4-68 years—see table 1). Ten of the 52 patients were symptomless but 35, 20, and five complained of dyspnoea, cough, and chest pain respectively. Physical examination revealed dullness and rales in 22, but no physical findings were noted in 22 of the patients. Cyanosis was noted in seven cases, clubbing in 12, and hepatomegaly in eight. Pectus excavatum and hypertrophic pulmonary osteoarthropathy were additional findings in two subjects.<sup>30,38</sup> The patients were from all the geographical regions of Turkey, though birthplaces were determined in only 23 cases.<sup>9,22,23,26</sup>

A family history was noted in 19 of the 39 patients. All the familial cases were siblings except for three who had a maternal aunt-nephew relationship. No family history was available in a quarter of the cases. Ten of the patients with a family history were female.


Nineteen of the patients had evidence of the disease on chest radiographs taken a mean of 3.9 years before the definitive diagnosis. All the radiographs showed the "sandstorm" appearances typical of pulmonary alveolar microlithiasis (figure). The initial diagnoses were sarcoidosis in one case, miliary tuberculosis in 13 cases and unknown in five. Only two with an initial diagnosis of miliary tuberculosis had acid fast bacilli in their sputum.

Although all of the 52 patients had a chest radiograph typical of pulmonary alveolar microlithiasis one patient also had bilateral apical pneumothoraces.<sup>9</sup> Bilateral diffuse calcification, pleural calcification, emphysematous bullae, and blebs were found in five other patients, who were examined by computed tomography.<sup>34,36,38,40,41</sup> Bone scintigraphy was also performed in seven patients and six had uptake of gallium in the lungs. Respiratory function tests were performed in 25 of the cases. The findings were normal in eight and the others all had a restrictive pattern. Arterial blood gas analysis in 15 patients

Table 1 Characteristics of pulmonary alveolar microlithiasis cases in Turkey

Reference	No of cases	Age (y) and sex	Family history	Diagnostic procedure
Eber <sup>8,24</sup>	1	20M	—	OLB
Tezok <sup>9</sup>	1	28M	—	CNB
Alpaslan <sup>10</sup>	1	20M	—	CNB
Üner <sup>11</sup>	1	16M	—	CNB
Vidinel <sup>12,13</sup>	3	24F, 30F, 42M	+, +, —	XR, CNB, OLB
Gerelioglu <sup>14</sup>	1	21M	—	OLB
Özdoğan <sup>15</sup>	1	22M	NA	OLB
Karasu <sup>16</sup>	1	18F	—	CNB
Özhan <sup>17</sup>	1	38F	—	CNB
Yücel <sup>18,24</sup>	1	21M	NA	OLB
Cakirca <sup>19,20,24</sup>	1	33M	—	CNB
Artvinli <sup>21</sup>	1	20M	—	OLB
Enacar <sup>22,23</sup>	6	10F, 12F, 15M 17M, 23F, 24M	+, +, +, —, +, +	XR, XR, XR XR, OLB, XR
Hacıhanefioglu <sup>24</sup>	4	20M, 21M, 40M 41M	NA, NA, NA NA	OLB, XR, OLB CNB
Akcan <sup>25</sup>	1	34F	NA	OLB
Kocuyigit <sup>26</sup>	3	40F, 44F, 50M	—, +, +	OLB, OLB, OLB
Yenel <sup>27</sup>	2	28F, 38M	+, +	OLB, OLB
Kiliçaslan <sup>28</sup>	2	20M, 23M	NA, NA	CNB, BAL
Dogan <sup>29</sup>	2	21M, 26M	NA, NA	OLB, OLB
Balkanli <sup>30</sup>	1	68M	—	BAL
Türktaş <sup>31</sup>	1	29M	NA	XR
Öztaskeent <sup>32</sup>	1	35M	—	XR
Kanra <sup>33</sup>	3	4M, 34F, 20M	+, +, +	XR, OLB, XR
Celikel <sup>34</sup>	1	40F	—	BAL, TBB
Yüksel <sup>35</sup>	2	?M, 35F	+, +	OLB, XR
Camsarı <sup>36</sup>	1	19M	—	BAL, TBB
Albayrak <sup>37</sup>	1	8M	—	OLB
Emri <sup>38</sup>	1	44F	—	OLB
Hazar <sup>39</sup>	4	30M, 46M, 17F, 8M	+, —, +, +	OLB, OLB, OLB OLB
Uçan <sup>40</sup>	1	21M	—	OLB
Kamali <sup>41</sup>	1	35M	NA	OLB

OLB—open lung biopsy; CNB—cutting needle biopsy; XR—chest radiograph; TBB—transbronchial biopsy; BAL—bronchoalveolar lavage; NA—unreported.



Chest radiograph showing typical sandstorm appearances of pulmonary alveolar microlithiasis

showed no hypoxaemia.

The diagnosis was made by open lung biopsy in 27 of the cases and by cutting needle biopsy, bronchoalveolar lavage, and transbronchial biopsy in nine, four, and four respectively. Radiological findings alone were considered to be sufficient in 12. Bronchoalveolar lavage fluid showed microliths in four of six patients, and transbronchial biopsy provided adequate material for histological examination in two of three patients.<sup>28 30 34-36 40</sup>

### Discussion

The mean age of our group was similar to that in the cases overall. Nine of our 52 patients were below the age of 18 (table 2).<sup>42 43</sup> The geographical distribution of the patients in Turkey was heterogeneous. Although pulmonary alveolar microlithiasis has been reported from all over the world, 52 (23%) of the total of 225 cases now reported come from Turkey.<sup>3 6 7 45</sup> In addition, some of the patients reported from foreign countries were ethnic Turks. We therefore believe that pulmonary alveolar microlithiasis is more prevalent among Turkish people.<sup>46 47</sup> Some of the patients reported from other countries had been exposed to certain dusts. None of our patients had an obvious history of such exposure.<sup>5-7 48 49</sup>

This review reports 52 cases of pulmonary alveolar microlithiasis from Turkey, 49 of which have previously been described only in

Turkish publications. Where family histories were available nearly half were positive for the disease, confirming previous reports that pulmonary alveolar microlithiasis may be a familial disease and suggesting that the gene is most common in Turkey.<sup>3-5 37 43 44 47 50</sup>

- 1 Friedrich N. Corpora amylacea in den Lungen. *Arch Pathol Anat* 1856;9:613-8.
- 2 Harbitz F. Extensive calcifications of the lungs as distinct disease. *Arch Intern Med* 1918;21:139-46.
- 3 Sosman MC, Dodd GD, Jones DW, Pillmore GU. The familial occurrence of pulmonary alveolar microlithiasis. *AJR* 1957;77:947-1012.
- 4 O'Neill RP, Cohn JE, Pellegrino ED. Pulmonary alveolar microlithiasis, a family study. *Ann Intern Med* 1967;67:957-67.
- 5 Vidinel I, Tellioglu E. Pulmoner alveolar mikrolityazis. *Tüberküloz ve Toraks* 1969;17:36-53.
- 6 Prakash UB, Barham SS, Rosenow EC, Brown ML, Payne WS. Pulmonary alveolar microlithiasis. *Mayo Clin Proc* 1983;58:290-300.
- 7 Fraser RG, Paré JAP, Paré PD, Fraser RS, Genereux GP. *Diagnosis of diseases of the chest*. Philadelphia: Saunders, 1991:2693-711.
- 8 Eber N, Sunay M, Uzunhasanoglu R. Pulmoner alveolar mikrolityazis. *Tüberküloz ve Toraks* 1962;10:243-50.
- 9 Tezok F, Balci K, Baris YI, Kurt C. Bir vaka münasebetiyle diffüz pulmoner alveolar mikrolityazis. *Tüberküloz ve Toraks* 1964;12:463-9.
- 10 Alpaslan I, Yagan R. Alveolar mikrolityazis. *Tüberküloz ve Toraks* 1966;14:463-8.
- 11 Üner E, Gürses H, Solu S, Menemenli N. Pulmoner alveolar mikrolithiasis. *Tüberküloz ve Toraks* 1966;14:469-72.
- 12 Vidinel I, Estürk Z, Tellioglu E, Önal A. Pulmoner alveolar mikrolityazis. *Ege Tıp Fak Der* 1967;6:399-403.
- 13 Vidinel I, Tellioglu E. Pulmoner alveolar mikrolityazis. In: *XIX Milli Türk Tıp Kongresi*. İstanbul: Celik Cilt, 1968:146-8.
- 14 Gerelioglu Y, Büyükcünal S, Sekban H, Akkaya A. Bir diffüz pulmoner alveolar mikrolithiasis vakası münasebetiyle. *Tüberküloz ve Toraks* 1967;15:123-6.
- 15 Özdoğan N, Aytug Z, Yazgan C. Pulmoner alveolar mikrolithiasis. In: *VIII Türk Tüberküloz Kongresi*. Ankara:Ongun Kardesler, 1968:388-94.
- 16 Karasu N, Öger O, Alper D, Özyardimci N. Diffüz pulmoner alveolar mikrolithiasis. *X Türk Tüberküloz Kongresi* Ankara:Ongun Kardesler, 1971:573-6.
- 17 Özhan EH, Karıncaoglu O. Pulmoner alveolar mikrolityazis vakası In: *X Türk Tüberküloz Kongresi*. Ankara:Ongun Kardesler, 1971:577-9.
- 18 Yücel EN, Isiklar H, Yurdakul S, Özdemir L. Pulmoner alveolar mikrolithiasis. In: *X Türk Tüberküloz Kongresi*. Ankara:Ongun Kardesler, 1971:580-2.
- 19 Cakirca R, Ekmekcioglu A, Aksüğü H. Pulmoner alveolar mikrolityazis. *Tüberküloz ve Toraks* 1971;19:296-300.
- 20 Cakirca R, Ekmekcioglu A, Aksüğü H. Pulmoner alveolar mikrolityazis. In: *X Türk Tüberküloz Kongresi*. Ankara:Ongun Kardesler, 1971:733-8.
- 21 Artvinli M, Özesmi M, Baris YI. Pulmoner alveolar mikrolityazis. *Hacettepe Tıp/Cerrahi Bül* 1972;5:407-11.
- 22 Enacar N, Karabiyikoglu G, Ekim N. Alveolar mikrolityazis. In: *XIII Türk Tüberküloz Kongresi*. İstanbul: Ongun Kardesler, 1977:522.
- 23 Enacar N, Yavuzer S, Sayli BS, Karabiyikoglu G, Ekim N. Pulmoner alveolar mikrolityazis. *Tüberküloz ve Toraks* 1978;26:186-96.
- 24 Hacıhanefioglu U, Uysal V. Alveolar mikrolithiasis (7 olgu üzerine patolojik inceleme). In: *III Ulusal Patoloji Kongresi*. İzmir:Bilgehan, 1978:462-70.
- 25 Akcan I, Coskun Y. Pulmoner alveolar mikrolithiasis. *Tüberküloz ve Toraks* 1982;30:37-9.
- 26 Kocyigit E, Gazioglu K, Cavdar T, Erben T. Ilginc özellikleri olan 3 alveolar mikrolithiasis vakası. *Solunum* 1983;8:276-83.
- 27 Yenel F, Küçükusta AR. Bir akciğer mikrolithiasisi olgusu. *Cerrahpaşa Tıp Fak Der* 1984;15:133-7.
- 28 Kılıçaslan Z, Küçükusta AR, Özbek I. Tanılabilen akciğer igne biyopsisi ve bronkoalveolar lavajla doğrulanmış iki akciğer mikrolithiasisi olgusu. *Ege Tıp Fak Der* 1985;24:1340-8.
- 29 Dogan R, Baltacıoglu T, Dölan M, Aksoy B. Pulmoner alveolar mikrolithiasis. *Tüberküloz ve Toraks* 1986;34:290-4.
- 30 Balkanlı S, Demir R, Patisroglu T, Özesmi M. Pulmoner alveolar mikrolithiasis tanısında bronkoalveolar lavaj. *Tüberküloz ve Toraks* 1987;35:212-6.
- 31 Türktaş H, Öztürk C, Güven M, Ugur P, Erzen C. Pulmonary alveolar microlithiasis with absence of technetium 99 m uptake of lungs. *Clin Nuclear Med* 1988;13:883-5.
- 32 Öztasent R, Demircioglu N, Özacar R. Idiopatik pulmoner alveolar mikrolithiasis. *İzmir Göğüs Hast Has*

Table 2 Age and sex distribution, presence of family history, and symptoms in patients reported worldwide and in our series

Reference	No of patients	Mean age (y)	Male (%)	Family history (%)	Symptomless (%)
Sosman <sup>3</sup>	45	31-50	46	56	70
O'Neill <sup>41</sup>	27	30-50	50	38	—
Kino <sup>43</sup>	51	5-9	—	61	—
Prakash <sup>6</sup>	8	23	62	0	62
Mascie-Taylor <sup>15</sup>	71	32	49	—	87
Volle <sup>46</sup>	40	—	57	52	—
Our series	52	27	65	48	19

- Der 1988;3:64-6.
- 33 Kanra G, Tanyol E, Göcmen A, et al. Pulmonary alveolar microlithiasis (a case report). *Turkish Journal of Pediatrics* 1988;30:61-7.
  - 34 Celikel T, Küllü S, Germen N, Bayik M, Oktay A. Confirmation of diagnosis of pulmonary microlithiasis by transbronchial biopsy: a case report. *Respir Care* 1989;34:989-92.
  - 35 Yüksel M, Öztekin I, Yılmaz G, Balkan E. Bir pulmoner alveolar mikrolitiazis olgusu. *Türk Patoloji Der* 1990;6:45-9.
  - 36 Camsari G, Kilic U, Cikirikcioglu S, Öztürk I, Ilker O, Hacıhanefioglu U. Pulmoner alveolar mikrolitiazis. *Türk Patoloji Der* 1990;6:7-10.
  - 37 Albayrak D, Gönül S, Tastepe I, Göcmen A. Ayin vakasi. *Yeni Tip Der* 1990;7:270-3.
  - 38 Emri S, Cöplü L, Selcuk TZ, Sahin AA, Baris YI. Hypertrophic pulmonary osteoarthropathy in a patient with pulmonary alveolar microlithiasis. *Thorax* 1991;46:145-6.
  - 39 Hazar A, Erbagci A, Caglar T, Yilmaz N, Erem AR. Mikrolitiazis tanisinda sintigrafinin degeri (4 vaka nedeniyle). In: *XIX Tüberküloz ve Göğüs Hastaliklari Kongresi*. Sivas: Dilek, 1991:304.
  - 40 Ucan ES, Keyf AI, Aydılek R, Yalcin Z, Kudu M, Sebit S, Ok U. 1962-1991 yıllari arasında Türkiye'de tani konulan alveolar mikrolitiazis olgulari. In: *Türkiye Solunum Arastirmalari Dernegi XIX Kongresi*. Bursa, Istanbul: Nazim Terzioğlu, 1991, 46.
  - 41 Kamali Ü, Yılmaz U, Utkaner G, Günel Ö, Budunelli T. Pulmoner alveolar mikrolitiaziste bilgisayarli tomografik ve patolojik bulgular (bir olgu nedeniyle). In: *Türkiye Solunum Arastirmalari Dernegi XIX Kongresi*. Bursa, Istanbul: Nazim Terzioğlu, 1991, 46.
  - 42 Thuraijasinam S, Dharmasena BD, Kasthuriratna T. Pulmonary alveolar microlithiasis. *Australias Radiol* 1975;19:175-80.
  - 43 Kino T, Kohara Y, Tsuji S. Pulmonary alveolar microlithiasis: a report of 2 young sisters. *Am Rev Respir Dis* 1972;105:105-10.
  - 44 O'Neill RP, Cohen JE, Pellegrino ED. Pulmonary alveolar microlithiasis a family study. *Ann Intern Med* 1967;67:957-67.
  - 45 Mascie-Taylor BH, Wardman AG, Madden CA, Page RL. A case of alveolar microlithiasis: observation over 22 years and recovery of material by lavage. *Thorax* 1985;40:952-3.
  - 46 Volle E, Kaufmann HJ. Pulmonary alveolar microlithiasis in pediatric patients: review of the world literature and two new observations. *Pediatr Radiol* 1987;17:439-42.
  - 47 Barnard NJ, Crocker PR, Blainey AD, Davies RJ, Ell SR, Levison DA. Pulmonary alveolar microlithiasis: a new approach. *Histopathology* 1987;11:639-45.
  - 48 Nouh MS. Is the desert lung syndrome (non occupational dust pneumoconiosis) a variant of pulmonary alveolar microlithiasis? *Respiration* 1989;55:122-6.
  - 49 Chinachoti N, Tangchai P. Pulmonary alveolar microlithiasis associated with the inhalation of snuff in Thailand. *Dis Chest* 1957;32:687-9.
  - 50 Garty I, Giladi N, Flatua E. Bone scintigraphy in two siblings with pulmonary alveolar microlithiasis. *Br J Radiol* 1985;58:763-6.